

Alpha Tubulin Recombinant antibody

Cat:B31003T

Company: HaoKeBio

Uniprot ID:P68363

Applications: IHC:1:1000-1:4000

Organism:Rabbit

IHC-Polymer:1:4000-1:16000

Species reactivity:Human Mouse Rat

IHC-TSA:1:5000-1:20000

Molecular Weight Calculation: 50 kDa

IF:1:50-1:200

Observed Molecular Weight: 50-55 kDa

WB:1:5000-1:50000

Background:

There are five tubulins in human cells: alpha, beta, gamma, delta, and epsilon. Tubulins are conserved across species. They form heterodimers, which multimerize to form a microtubule filament. An alpha and beta tubulin heterodimer is the basic structural unit of microtubules. The heterodimer does not come apart once formed. The alpha and beta tubulins, which are each about 55 kDa MW, are homologous but not identical. Alpha tubulin is useful for scientists across fields as an internal control due to its high, ubiquitous expression pattern.

Protein full name:

tubulin, alpha 1b

Synonyms:

TUBA1B, Tubulin alpha 1B chain, 1G7, Alpha-tubulin ubiquitous, EC:3.6.5.-

Immunogen:

Recombinant protein

Isotype:

IgG

Subcellular location:

Cytoskeleton

Purity:

Affinity purification

Form:

Liquid

Storage Buffer:

PBS with 0.02% sodium azide, 100 µg/ml BSA and 50% glycerol.


Storage:

Store at -20 °C for one year.

Experimental procedure:

Antigen retrieval: Citrate buffer (pH 6.0), Medium high heat for 8 minutes, stop for 7 minutes, medium high heat for 8 minutes. Incubate antibody, 4°C overnight. Secondary antibody: Poly-HRP Goat Anti-Rabbit & Mouse Universal Secondary Antibody, RT, 1h.

Images:



Dilution of 1:50000 incubated at room temperature for 1.5 hours.

Source of Reagents:

发表[中文论文]请标注:Alpha Tubulin(B31003T)由杭州浩克生物技术有限公司提供;

发表[英文论文]请标注:Alpha Tubulin(B31003T) were kindly provided by Hangzhou Haoke Biotechnology Co., Ltd.